

Assessing essential oil composition in *Cinnamomum cassia* leaves from different regions of China using GC-MS and FTIR spectroscopy

YAN HUANG¹, YUJIE LIU¹, HUIPING TAN¹, YANRONG CHENG¹, KUNYANG TAO¹,
DINGZE GU¹, HUAIZU CAI¹, CHENGJIE LI¹, KAIYI GUO¹, CHENG WU¹,
HONG WU^{1,2*}, YANQUN LI^{1,2*}

¹Guangdong Key Laboratory for Innovative Development and Utilization of Forest Plant Germplasm,
South China Agricultural University, Guangzhou, China

²Medicinal Plants Research Center, South China Agricultural University, Guangzhou, China

*Corresponding authors: wh@scau.edu.cn; liyanqun@scau.edu.cn

Yan Huang and Yujie Liu contributed equally to this work.

Citation: Huang Y., Liu Y., Tan H., Cheng Y., Tao K., Gu D., Cai H., Li C., Guo K., Wu C., Wu H., Li Y. (2024): Assessing essential oil composition in *Cinnamomum cassia* leaves from different regions of China using GC-MS and FTIR spectroscopy. Czech J. Food Sci., 42: 141–152.

The authors are fully responsible for both the content and the formal aspects of the Electronic Supplementary Material. No editorial adjustments were made.

Electronic Supplementary Material

Supplementary Figures S1–S2




Figure S1. Gas chromatography-mass spectrometry profiles (A–C) of the essential oils from the leaves of *Cinnamomum cassia* and (D) of *trans*-cinnamaldehyde standard sample




Figure S2. Full Fourier transform infrared spectra of essential oils from nine samples