

Biogenic Amine Content in Sterilised and Pasteurised Long-Term Stored Processed Cheese

TOMÁŠ KOMPRDA¹, KATEŘINA NOVICKÁ¹, LIBOR KALHOTKA¹ and DANA SMĚLÁ²

¹Department of Food Technology, Faculty of Agronomy, Mendel University of Agriculture and Forestry Brno, Brno, Czech Republic; ²Czech Agriculture and Food Inspection Authority, Brno, Czech Republic

Abstract

KOMPRDA T., NOVICKÁ K., KALHOTKA L., SMĚLÁ D. (2005): **Biogenic amine content in sterilised and pasteurised long-term stored processed cheese.** Czech J. Food Sci., **23**: 209–216.

The contents of eight biogenic amines (tryptamine, phenylethylamine, histamine, tyramine, putrescine, cadaverine, spermidine and spermine) were determined in samples of processed cheese, either pasteurised (P) or sterilised (S), and consequently stored for either 22 or 57 weeks at the temperatures of 8°C and 22°C, respectively. Tyramine was quantitatively the most important biogenic amine (1.3–29.3 mg/kg); the highest level ($P < 0.05$) was found in P cheese stored for 22 weeks after processing at 8°C (P1). Regardless of tyramine, and with the exception of putrescine in S cheese stored for 57 weeks (2.2 mg/kg), the content of no biogenic amine exceeded 2 mg/kg in any tested sample. Significantly higher (by five to six orders of magnitude; $P < 0.05$) counts of total aerobic and facultative anaerobic microorganisms in P1 sample in comparison with all other samples were indicative of the cover leakage. No lactic acid bacteria were detected in any sample, the counts of coliforms and total anaerobes, respectively, were negligible and did not differ ($P > 0.05$) between samples. No sample of the tested long-term stored processed cheese was considered unsafe from the toxicological viewpoint.

Keywords: tyramine; polyamines; spoilage bacteria; food safety

Cheese, considered on the one hand healthy and recommendable food, is on the other hand one of the most important suppliers of biogenic amines (BA), the low-molecular organic bases formed mainly by decarboxylation of amino acids (SILLA-SANTOS 1996), into the human organism. This usually poses no problem with healthy people, but with allergic individuals or people administered monoamine oxidase (MAO) inhibitors, negative health effects after the intake of relatively low levels of BA (e.g. 6 mg of tyramine; SILLA-SANTOS 1996) can be expected. Apart from the vasoactive amines tyramine (hypertension, headache)

or histamine (hypotension, headache, flushing, abdominal cramps; STRATTON *et al.* 1991), growing attention is paid at present to the polyamines spermidine and spermine, including diamines putrescine and cadaverine, due to their ability to form stable carcinogenic N-nitroso compounds and to enhance the growth of chemically induced aberrant crypt foci in the intestine (PAULSEN *et al.* 1997). Polyamines, both formed endogenously and taken from the diet, can be involved in the tumor development (KALAČ & KRAUSOVÁ 2005).

Although BA content has been evaluated sufficiently in ripening cheeses (PETRIDIS & STEIN-

HART 1996a, b), the data regarding processed cheese are very scarce in the available literature (EL-SAYED 1996).

The two objectives of the present experiment were as follows. Firstly, within the scope of quality evaluation of the processed cheese produced for the military purposes of the long-term storage, to compare the contents of biogenic amines in the samples either sterilised or pasteurised after processing and subsequently long-term stored at the refrigerator and at room temperature, respectively. Secondly, in connection with the former task, to evaluate the suitability of the high-pressure liquid chromatography method for the determination of biogenic amines in processed cheese, i.e. the matrix with presumably very low biogenic amine content.

MATERIALS AND METHODS

Cheese samples. The cheese samples were produced for the military purposes of the long-term storage ("military academy" samples). On delivery to our laboratory, the cheese samples were declared to have been produced within one batch consisting of the same ingredients: blocks of Edam cheese (45% fat in dry matter), Moravian block cheese, butter, curd, water, and an emulsifying agent. All ingredients were put into the processed cheese kettle equipped with a steam-heated shell and a direct steam injection. The mixture was processed 10 min at the temperature of 85°C under stirring. The processed matter was filled into aluminum boxes in the amounts of 200 g and the lids were fused. The set of the samples of the processed cheese was divided into two parts. One part was cooled to the temperature of 10°C, this cheese was designated as pasteurised processed cheese – P (that is the cheese treated only by the temperature 85°C for 10 min). The second part of the samples was put into a laboratory autoclave (Labo Autoclave Sanyo, model MLS 3780), sterilised 20 min under the pressure of 0.25 MPa at the temperature of 117°C, and subsequently cooled. These samples were designated as sterilised processed cheese – S. The samples of both pasteurised and sterilised processed cheese were stored either 22 or 57 weeks, the pasteurised cheese at the temperature of 8°C, the sterilised cheese at the temperature of either 8°C or 25°C (Table 1).

Biogenic amine determination. 10 g (\pm 1 mg) of the sample was weighed into an 85 ml test tube,

Table 1. Identification of the analysed samples of processed cheese

Sample designation	<i>n</i>	Heat treatment	Storage temperature (°C)	Storage time (weeks)
S1	2	sterilisation	8	22
S2	4		8	57
S3	4		25	57
P1	2	pasteurisation	8	22
P2	4		8	57

n – number of tested samples; each sample was measured in duplicate

0.5 ml of an internal standard (1,7-diaminoheptane; concentration 1 mg/ml) was added and the sample was extracted for two minutes with 20 ml of 0.1M HCl using the disintegrator Heidolph Diax 900 (Germany). The suspension was centrifuged at $755 \times g$ for 10 min at 4°C. The supernatant was filtered through paper filter and the solid residue was repeatedly extracted as above. The combined extracts were made up to 50 ml with deionised water and filtered through a disposable nylon membrane filter (13 mm, 0.45 μ m).

An aliquot of the extract was derivatised with dansyl chloride (5-dimethylaminonaphthalene-1-sulfonyl chloride, DCl) as follows: 1 ml of the extract (or standard) was mixed with 0.5 ml of saturated Na₂CO₃, 1 ml of DCl was added (Sigma-Aldrich, Germany; the agent was prepared by dissolving of 5 mg of DCl in 1 ml of acetone), and the mixture was shaken for 1 min (MS2 Minishaker IKA Werke, Germany). Derivatisation proceeded for 1 h in the dark at 40°C. After derivatisation, 250 μ l of ammonia was added to remove excess DCl. After 30 min, the amine derivatives were extracted by diethylether (3 \times 1 ml), the organic phase was evaporated to dryness under nitrogen, and the solid residue was dissolved in 0.5 ml of acetonitrile (ACN). The solution was filtered through the nylon membrane filter 0.45 μ m and injected onto the chromatographic column.

Biogenic amines were separated using a liquid chromatograph HP 1100 (Agilent Technologies, Waldbronn, Germany) consisting of the quaternary pump (G1311A), vacuum degasser (G1322A), automatic sampler (G1313A), UV/VIS detector with the variable wave-length (G1314A) and a fluorescence detector (G1321A). The separation

after DCl derivatisation was carried out by gradient elution with H₂O/ACN (time 0–23 min: H₂O 35–0%, ACN 65–100%) on the Zorbax Eclipse XDB C18 column (150 mm × 4.6 mm, particle size 5 µm) with the guard column Meta Guard ODS-2 (30 mm × 4.6 mm, particle size 5 µm) at the flow rate of 0.8 ml/min using photometric UV/VIS detector at 254 nm.

Regarding several steps of the sample preparation, the concentration of a given biogenic amine in the sample (c_x) was corrected based on the method of internal standard as follows:

$$c_x = c_{IS} \times A_x / A_{IS} \times RF_x$$

where: c_{IS} – concentration of an internal standard (in mg/kg)

A_{IS} – peak area of the internal standard (in area units)

A_x – peak area of the biogenic amine (in area units)

RF_x – response factor of the amine

$$RF_x = c_{xr} / c_{ISr} \times A_{ISr} / A_{xr}$$

where: c_{xr} – concentration of biogenic amine in the reference sample (mg/kg)

c_{ISr} – concentration of an internal standard added to the reference sample (mg/kg)

A_{ISr} – peak area of the internal standard in the reference sample (in area units)

A_{xr} – peak area of the biogenic amine in the reference sample (in area units)

The separated amines (histamine, tyramine, tryptamine, putrescine, 2-phenylethylamine, cadaverine, spermidine, spermine) were identified by comparison of the retention times of the standards (used as the respective hydrochlorides, Sigma-Aldrich, Germany). Their concentrations after DCl derivatisation were expressed in mg/kg of the original (fresh) matter.

The repeatability of the determination and the recoveries of particular biogenic amines were estimated with a cheese sample different from the “military academy” samples used in the experiment proper, with no strictly defined origin, but with a supposedly similar (that is relatively low) biogenic amine content (a “standard” sample). Five extracts of this sample (prepared as above-mentioned) after derivatisation with DCl were used for the repeatability (expressed as a relative standard deviation, RSD in %) determination of the entire analytical procedure (method). Ten times injected mixture of the BA standards after DCl derivatisation was used for characterising

the repeatability of the chromatographic system (instrument). The recoveries were evaluated using the “standard” sample spiked with the mixture of the BA standards on the concentration level 1 and 2 mg/kg, respectively, measured five times. The recovery (R) was calculated as

$$\% R = [(CF - CU) / CA] \times 100$$

where: CU – concentration in the original sample

CA – concentration of the analyte added

CF – concentration in the spiked sample

The concentrations of the particular amines in the “military academy” samples of long-term stored processed cheese were not corrected for the respective recovery.

Microbiological analysis. 20 g of cheese were taken aseptically from the sample, homogenised in the stomacher with 180 ml of distilled water, and serial decimal dilutions were prepared. The following groups of microorganisms were determined: total counts of aerobic and facultative anaerobic mesophilic microorganisms (TAC) on Plate Count Agar (PCA; Biokar Diagnostics, France) after 72 h at 30°C; lactic acid bacteria (LAB) on De Man-Rogosa-Sharpe medium (MRS; Biokar) after 72 h at 37°C; coliforms on Violet Red Bile Agar (Biokar) after 72 h at 37°C; total anaerobic counts (TAN) on PCA (Biokar) after 120 h at 30°C (1 ml of the sample was overlaid in the test tube with the 10 cm agar column); yeasts + moulds on glucose-yeast extract-chloramphenicol agar (Noack, Vienna, Austria) after 120 h at 25°C.

Statistical evaluation. One-way classification of the variance-ratio test, including Duncan’s multiple range test, was used for the calculation of the difference between biogenic amine contents in the samples using the Unistat package, version 4.53 (Unistat, Ltd., London).

RESULTS

The data characterising the repeatability (expressed as a relative standard deviation, RSD) of the chromatographic system (instrument) and of the entire analytical procedure (method), and the recovery of each biogenic amine spiked at two levels, are presented in Table 2. Relative standard deviation calculated from the measured data regarding five extracts of the “standard” cheese sample (different from the samples shown in Table 1) was within the range of 1.9% for spermidine to

Table 2. Repeatability (expressed as a relative standard deviation, RSD in %) of the biogenic amine determination, and recovery of the biogenic amines determined in the cheese samples

Biogenic amine	Instrument ¹ (RSD, %)	Method ² (RSD, %)	Recovery ³ (%)	
			1 mg/kg	2 mg/kg
Tryptamine	0.3	ND ⁴	75.1	94.9
Phenylethylamine	0.4	6.8	78.4	90.3
Histamine	0.3	5.8	115.4	93.6
Tyramine	1.6	5.0	59.8	56.1
Putrescine	0.3	3.0	65.1	57.2
Cadaverine	0.3	2.9	74.6	71.3
Spermidine	0.3	1.9	63.3	52.6
Spermine	0.2	5.4	66.6	50.2

¹mixture of the biogenic amine standards after derivatisation was injected ten times; ²five extracts of the same cheese sample after derivatisation was used; ³cheese sample spiked with the mixture of biogenic amine standards on the concentration level 1 and 2 mg/kg, respectively, was measured five times; ⁴not detected

6.8% for phenylethylamine. The recovery of the biogenic amines determined in the “standard” cheese sample spiked with the mixture of BA standards at the concentration level 2 mg/kg was in the range of 50.2% for spermine to 94.9% for tryptamine.

Biogenic amine contents in the samples of long-term stored processed cheese from military academy are shown in Table 3. Quantitatively most important biogenic amine in all tested samples was tyramine. The tyramine content in pasteurised cheese stored for 22 weeks after processing (P1)

was significantly higher ($P < 0.05$) in comparison with the sterilised cheese stored for the same time interval (S1) and also in comparison with all other tested samples. On the other hand, the tyramine content in sterilised cheese stored for 22 weeks at the temperature of 8°C was lower ($P < 0.05$) than in the sterilised samples stored for 57 weeks both at the temperature of 8°C and at the temperature of 25°C.


Regardless of tyramine, and with the exception of putrescine in S2 (cheese sterilised after processing

Table 3. Biogenic amine contents in the samples of the long-term stored processed cheese

Biogenic amine	Sample (mg/kg of the cheese; mean \pm standard error of the mean)				
	P1 ¹	S1 ²	P2 ³	S2 ⁴	S3 ⁵
Tryptamine	0.7 ^A \pm 0.10	0.0 ^A \pm 0.00	0.2 ^A \pm 0.10	0.2 ^A \pm 0.06	0.8 ^A \pm 0.75
Phenylethylamine	1.8 ^A \pm 1.40	0.2 ^B \pm 0.00	0.3 ^A \pm 0.03	0.7 ^{A,B} \pm 0.14	0.3 ^A \pm 0.94
Histamine	0.7 ^A \pm 0.35	0.0 ^A \pm 0.00	0.3 ^A \pm 0.21	0.7 ^A \pm 0.14	0.4 ^A \pm 0.23
Tyramine	29.3 ^C \pm 6.90	1.3 ^A \pm 0.05	6.3 ^{A,B} \pm 0.56	9.3 ^B \pm 0.45	7.7 ^B \pm 0.19
Putrescine	1.0 ^A \pm 0.35	0.3 ^A \pm 0.05	1.9 ^A \pm 0.33	2.2 ^A \pm 0.61	1.6 ^A \pm 0.60
Cadaverine	0.7 ^{A,B} \pm 0.05	0.5 ^A \pm 0.05	1.3 ^{A,B} \pm 0.29	1.2 ^B \pm 0.12	1.0 ^{A,B} \pm 0.13
Spermidine	0.2 ^B \pm 0.00	0.2 ^B \pm 0.00	0.1 ^A \pm 0.00	0.2 ^B \pm 0.03	0.1 ^{A,B} \pm 0.03
Spermine	1.0 ^B \pm 0.30	0.0 ^A \pm 0.00	0.1 ^A \pm 0.13	0.2 ^A \pm 0.18	0.3 ^A \pm 0.13

¹pasteurised, stored 22 weeks at 8°C; ²sterilised, stored 22 weeks at 8°C; ³pasteurised, stored 57 weeks at 8°C; ⁴sterilised, stored 57 weeks at 8°C; ⁵sterilised, stored 57 weeks at 25°C;

^{A,B,C} means with different superscripts in lines differ significantly ($P < 0.05$)



A,B,C means with different superscripts within the particular group differ significantly ($P < 0.05$)

Figure 1. Sum of all determined biogenic amines (tryptamine, phenylethylamine, histamine, tyramine, putrescine, cadaverine, spermidine, spermine) and sum of polyamines (putrescine, cadaverine, spermidine, spermine) in the samples of long-term stored processed cheese

and stored for 57 weeks thereafter), the content of no biogenic amine exceeded 2 mg/kg in any sample tested.

The highest ($P < 0.05$) content of all biogenic amines (calculated as a sum of all determined BA) was found in pasteurised cheese stored for 22 weeks at 8°C (P1; Figure 1). On the other hand, the sum of all BAs in the sterilised cheese stored for the same time interval at the same temperature (22 weeks at 8°C; S1) was lower ($P < 0.05$) not only in comparison with P1 sample, but also in comparison with both sterilised samples stored for 57 weeks at both temperatures (8°C and 25°C, respectively). Also the sum of polyamines in S1 cheese was lower ($P < 0.05$) in comparison with the sterilised sample stored for 57 weeks both at the refrigerating temperature and at the room temperature. However, generally speaking, the sum of polyamines was very low in all cheeses and did not exceed 2.5 mg/kg in any sample.

As far as microbiological analysis is concerned, the total aerobic counts in P1 sample were 1.3×10^6 cfu (colony-forming units) per g, which was a higher ($P < 0.05$) value in comparison with TAC in all the other samples of the processed cheese (the range 2.5×10^0 to 6.5×10^1 cfu/g; the differences were insignificant, $P > 0.05$). The analyses of lactic acid bacteria and yeasts and moulds, respectively, gave negative results in all tested samples of the

processed cheese in the present experiment. Coliforms were detected (in negligible amounts, 11 to 28 cfu/g) only in samples stored for 57 weeks. The counts of anaerobic bacteria were also very low in all analysed samples (the range 11 to 35 cfu/g). No differences ($P > 0.05$) between the samples of the long-term stored processed cheese were found regarding the content of coliforms and TAN, respectively (detection of LAB and yeasts + moulds, as mentioned above, was negative in all cases).

DISCUSSION

Because only strictly limited number of two-hundred g samples of the long-term stored processed cheese was available for both chemical and microbiological analyses, the qualimetric determinations were carried out with different cheese samples than those used in the experiment proper, but with supposedly similar contents of biogenic amines. Our results regarding the repeatability of the chromatographic system and analytical procedure, including BA recoveries (Table 2), are fully comparable with, in many cases even better than, the data of MORET and CONTE (1996), who reported the repeatability in the range of 3.4% for tyramine to 12.5% for spermidine when analysing the biogenic amine content in Grana Padano cheese. As far as the recoveries are concerned,

our results are not worse (often better) both in a comparison with the data of MORET and CONTE (1996) and VALE and GLÓRIA (1997) regarding biogenic amines in Grana Padano cheese (2.5% for putrescine to 79% for spermidine) and Parmesan cheese (71.4% for spermidine to 95.4% for histamine), respectively.

Regarding one of the objectives of the present study, i.e. testing the surmised effect of the processing temperature on the microbial status and, consequently, on the BA content, the higher tyramine concentration in P1 than in S1 sample (pasteurised vs. sterilised cheese stored 22 weeks) was contrary to the expectations. There are at least three acceptable explanations for the conspicuously different ($P < 0.05$) tyramine content in P1 as compared not only to S1 but also to all other samples.

Firstly, the obvious reason for the presence of biogenic amines in processed cheese is the transfer of these substances from the ripening cheese used as a raw material in the processed cheese production. Based on the data of EL-SAYED (1996), an effect of high temperature on biogenic amine content already present in the raw material (ripening cheese) is improbable. One of the possible explanations of the above difference between P1 and S1 samples in the present study is therefore the use of different raw materials for the production of processed cheese designated as pasteurised and sterilised, respectively (Table 1). In other words, the samples of the processed cheese, which were delivered to our laboratory, could have originated not from a single batch as declared, but from two (or even more) batches.

Another possibility, still presuming the samples originated from a single batch, is the use of raw materials (Edam cheese and Moravian block) with varying original tyramine contents, which were not uniformly distributed in the resulting cheese during processing.

Biogenic amines (with the exception of spermidine and spermine) are usually formed in cheese by an action of microbial decarboxylases. Therefore, thirdly, the surprisingly higher ($P < 0.05$) tyramine content in pasteurised cheese stored for 22 weeks (P1) as compared even to sterilised cheese stored 57 weeks at the room temperature (S3; Table 3) could be most probably indicative of the secondary microbial contamination as a consequence of the cover leakage. This possibility was supported by the data of the microbiological analysis. Notwithstanding either zero or negligible (and not

different, $P > 0.05$) counts of LAB, TAN, coliforms, and yeasts + moulds, respectively, significantly higher ($P < 0.05$) counts of total aerobes were found in P1 sample in comparison with TAC in all the other samples of the processed cheese. It is generally accepted that the counts of microorganisms, usually analysed in foods, do not often correlate with the biogenic amine content, due to the fact that the amine production is not the matter of particular microbial groups, genera or species, but rather the matter of particular strains within the given species (BOVER-CID *et al.* 2000). However, the above difference of five to six orders of magnitude in the TAC between P1 sample and all other samples, together with nearly five times as high tyramine content in P1 in comparison with an average value of four other samples in the present experiment, is sufficiently persuasive of the secondary contamination by microorganisms, including some strains (not specified in detail in the present experiment) possessing decarboxylase activity.

As far as both pasteurised samples (P1 vs. P2) are concerned, no reasonable conclusion was possible to draw regarding the storage time (these samples differed only in this parameter), because both the tyramine content (Table 3) and the content of total BA (Figure 1) were lower ($P < 0.05$) in the sample stored by 35 weeks longer. Contrary to this result, it would be logical to presume an increase of BA content with increasing storage time (VALSAMAKI *et al.* 2000). Biogenic amine degradation in the food matrix is possible (LEUSCHNER *et al.* 1998), but in the present experiment improbable (although it is true that the microorganisms with a known potential to degrade biogenic amines, especially histamine and tyramine, e.g. some strains of *Brevibacterium linens*, *Staphylococcus carnosus*, *Geotrichum candidum*, *Micrococcus varians*, were not analysed in the present experiment). It could be concluded that any possible effect of the storage time was masked by the above-mentioned uncontrolled factors.

A similar conclusion follows from the assessment of the surmised effect of the heat treatment regarding the comparison of the samples P2 and S2 (pasteurised and sterilised samples, respectively, stored at the same temperature for the same time period). Though not statistically significant ($P > 0.05$), both the tyramine content (Table 1) and the content of total BA (Figure 1) in the sterilised samples (S2) tended to be higher

in comparison with the pasteurised samples (P2). This, again, is not logical, if the heat treatment is considered the only important controlling factor (more thorough elimination of microorganisms by sterilisation). Because microbial analysis did not provide any guidance in this case (no differences in the microbial traits between the samples P2 and S2 were found, and therefore no secondary contamination could be supposed), the most probable explanation is the production of the P2 and S2 samples not within one batch (as declared), but within two different batches processed from the raw materials (i.e. ripening cheeses) with different original BA contents.

Contrary to the effect of the storage time in the pasteurised samples (see above), the higher ($P < 0.05$) tyramine content and the higher ($P < 0.05$) content of the sum of BA in the sterilised samples S2 and S3 in comparison with the sample S1 (Table 3, Figure 1) could be explained by a longer storage interval (57 weeks in the case of S2 and S3, 22 weeks in the case of the sample S1). However, also this explanation should be taken with caution, because the increase of BA content with increasing storage time is relevant only for the ripening cheese (VALSAMAKI *et al.* 2000). In properly produced and stored processed cheese, no substantial increase of any microorganisms possessing decarboxylase activity (in fact of any microorganisms whatsoever) should be presumed. It is questionable if the amounts of coliform microorganisms found in the samples stored for 57 weeks (11–28 cfu/g) in the present experiment could be sufficient to cause the observed differences in BA content.

The samples of processed cheese evaluated in the present study were analysed on request of a military academy. From the methodological viewpoint, when the samples were delivered to our laboratory, the only presumed variability factors which were declared, were the processing temperature, the storage time, and the storage temperature. However, as follows from the results, some other factors influencing biogenic amine content in the analysed samples were probably not sufficiently controlled or the conditions of the sample preparation and storage were not properly kept. Therefore no conclusions regarding the effect of the processing temperature, storage temperature or storage time period on the biogenic amine content in the processed cheese were possible in the present experiment.

The only possible conclusion is a general statement that none of the tested samples, including P1, would present health hazard from the toxicological viewpoint. Toxicological level of tyramine, the most toxic biogenic amine (a potent vasoactive substance), is above 100 mg/kg of the consumed food (SILLA-SANTOS 1996). Similarly, the sum of all biogenic amines in the tested samples in the present experiment, including P1 (Figure 1), was well under the safety limit of 900 mg/kg for the sum of tyramine + histamine + putrescine + cadaverine according to VALSAMAKI *et al.* (2000). One-hundred gram portion of the P1 cheese would be considered safe even in patients receiving non-selective MAO inhibitors, where the toxic dose was suggested to be 6 mg of tyramine (NOVELLA-RODRÍGUES *et al.* 2003).

Moreover, biogenic amine content in the cheese samples analysed in the present experiment was substantially lower not only in comparison with usual levels being found in ripening cheeses (e.g. histidine, tyramine, cadaverine and putrescine content in Swiss-type cheese stored for 5 months at 5°C was 250, 320, 170 and 160 mg/kg, respectively, in the experiment of PETRIDIS and STEINHART (1996b)), but even in comparison with the average values within the set of different kinds of processed cheese reported by EL-SAYED (1996): tyramine 58.7 mg/kg, spermine 56.0, putrescine 21.1 mg/kg.

The use of milk of a good hygienic quality and, consequently, the use of the ripening cheese in relatively low ripening grade as a raw material for the production of the processed cheese analysed in the present experiment, can be implied from the above comparisons and also from a very low content of polyamines spermidine and spermine in the final product (Table 3). Spermidine and spermine are not formed by the microbial action in the cheese but pass from the milk to the product (NOVELLA-RODRÍGUES *et al.* 2003).

At present, polyamines putrescine, spermidine and spermine are intensively studied due to their involvement in the process of carcinogenesis (KALAČ & KRAUSOVÁ 2005). It is known (WOLTER *et al.* 2004), that the effect of some cytostatics (inhibition of one of the key enzymes of the polyamine synthesis in the mammalian organism, ornithine decarboxylase) can be reversed by an increased polyamine uptake from food. Processed cheese analysed in the present experiment can be considered safe from this viewpoint (very low polyamine levels are apparent from Figure 1).

CONCLUSIONS

The results of the present study demonstrated the suitability of the high-pressure liquid chromatography method, including dansyl chloride derivatisation and UV detection, for the determination of biogenic amines in processed cheese, a matrix with a very low levels of these toxicologically interesting substances.

On the other hand, no conclusion regarding a surmised effect of the processing temperature, storage temperature, or storage time period on the biogenic amine content in the processed cheese analysed in this experiment was possible. Two likely reasons were considered. Firstly, some other factors apart from the above-mentioned (processing temperature, storage temperature, storage time interval) were not controlled, e.g. not the same raw materials were used for the production of all samples of the processed cheese. Secondly, though not visibly apparent, some cover damage of one of the samples was supposed based on the microbial analysis and the outlying results regarding tyramine content in this sample. It could only be concluded that, despite the above shortcomings, none of the tested samples would present health hazard from the toxicological viewpoint.

Acknowledgement. The authors also wish to thank Dr. FRANTIŠEK BUŇKA, Tomáš Baťa University Zlín, for providing the samples of processed cheese.

References

- BOVER-CID S., IZQUIERDO-PULIDO M., VIDAL-CAROU M.C. (2000): Influence of hygienic quality of raw materials on biogenic amine production during ripening and storage of dry fermented sausage. *Journal of Food Protection*, **63**: 1544–1550.
- EL-SAYED M.M. (1996): Biogenic amines in processed cheese available in Egypt. *International Dairy Journal*, **6**: 1079–1086.
- KALAČ P., KRAUSOVÁ P. (2005): A review of dietary polyamines: Formation, implications for growth and health and occurrence in foods. *Food Chemistry*, **90**: 219–230.
- LEUSCHNER R., HEIDEL M., HAMMES W.P. (1998): Histamine and tyramine degradation by food fermenting microorganisms. *International Journal of Food Microbiology*, **39**: 1–10.
- MORET S., CONTE L.S. (1996): High-performance liquid chromatography evaluation of biogenic amines in food. An analysis of different methods of sample preparation in relation to food characteristics. *Journal of Chromatography A*, **729**: 363–369.
- NOVELLA-RODRÍGUES S., VECIANA-NOGUÉS M.T., IZQUERDO-PULIDO M., VIDAL-CAROU M.C. (2003): Distribution of biogenic amines and polyamines in cheese. *Journal of Food Science*, **68**: 750–755.
- PAULSEN J.E., REISTAD R., ELIASSEN K.A., SJAASTAD O.V., ALEXANDER J. (1997): Dietary polyamines promote the growth of azoxymethane-induced aberrant crypt foci in rat colon. *Carcinogenesis*, **18**: 1871–1875.
- PETRIDIS K.D., STEINHART H. (1996a): Biogene Amine in der Hartkäseproduktion: I. Einfluß verschiedener Parameter auf den Amingehalt im Endprodukt am Beispiel von Emmentaler Käse. *Deutsche Lebensmittel-Rundschau*, **92**: 114–120.
- PETRIDIS K.D., STEINHART H. (1996b): Biogene Amine in der Hartkäseproduktion: II. Stufenkontroll-Studie einer standardisierten Emmentalerkäse-Produktion. *Deutsche Lebensmittel-Rundschau*, **92**: 142–146.
- SILLA-SANTOS M.H. (1996): Biogenic amines: their importance in foods. *International Journal of Food Microbiology*, **29**: 213–231.
- STRATTON J.E., HUTKINS R.V., TAYLOR S.L. (1991): Biogenic amines in cheese and other fermented foods. A review. *Journal of Food Protection*, **54**: 460–470.
- VALE S., GLÓRIA M.B. (1997): Determination of biogenic amines in cheese. *Journal AOAC International*, **80**: 1006–1012.
- VALSAMAKI K., MICHAELIDOU A., POLYCHRONIADOU A. (2000): Biogenic amine production in Feta cheese. *Food Chemistry*, **71**: 259–266.
- WOLTER F., ULRICH S., STEIN J. (2004): Molecular mechanisms of the chemopreventive effects of resveratrol and its analogs in colorectal cancer: Key role of polyamines? *Journal of Nutrition*, **134**: 3219–3222.

Received for publication April 15, 2005
Accepted after corrections June 9, 2005

Corresponding author:

Prof. MVDr. Ing. TOMÁŠ KOMPRDA, CSc., Mendelova zemědělská a lesnická univerzita v Brně, Agronomická fakulta, Ústav technologie potravin, Zemědělská 1, 613 00 Brno, Česká republika
tel.: + 420 545 133 261, fax: + 420 545 133 190, e-mail: komprda@mendelu.cz
